

Use of Protein: creatinine ratio in a random spot urine sample for predicting significant proteinuria in diabetes mellitus

BK Yadav,¹ S Adhikari,² P Gyawali,¹ R Shrestha,¹ B Poudel¹ and M Khanal¹

¹Department of Biochemistry, Tribhuvan University Teaching Hospital and ²National College for advance learning, Purbanchal University, Kathmandu, Nepal

Corresponding author: Mr. Binod Kumar Yadav, Lecturer; IOM, TUTH, Kathmandu, Nepal; e-mail: binod3aug@gmail.com

ABSTRACT

Present study was undertaken during a period of 6 months (September 2008-February 2009) to see an correlation of 24 hours urine protein estimation with random spot protein-creatinine (P:C) ratio among a diabetic patients. The study comprised of 144 patients aged 30-70 years, recruited from Kantipur hospital, Kathmandu. The 24-hr urine sample was collected, followed by spot random urine sample. Both samples were analyzed for protein and creatinine excretion. An informed consent was taken from all participants. Sixteen inadequately collected urine samples as defined by (predicted creatinine - measured creatinine)/predicted creatinine >0.2 were excluded from analysis. The Spearman's rank correlation between the spot urine P:C ratio and 24-hr total protein were performed by the Statistical Package for Social Service. At the P:C ratio cutoff of 0.15 and reference method (24-hr urine protein) cutoff of 150 mg/day, the correlation coefficient was found to be 0.892 (p<0.001). The area under ROC curve at different cutoffs was 0.88 at 95.0% CI. The sensitivity and specificity of the P:C ratio to detect significant proteinuria at the cutoff of 0.15 are 96.6% and 74.4%. So the P:C ratio can predict significant proteinuria in diabetic subjects, avoiding the inconvenient 24-hr urine collection but the cutoff should be carefully selected for different patients group under different laboratory procedures and settings.

Keywords: 24-hr urinary protein, Protein: creatinine ratio, Proteinuria, ROC curve.

INTRODUCTION

Nephropathy is the major health problem in patients with diabetes mellitus (DM). The natural history of diabetic nephropathy has generally viewed as a descending path from normoalbuminuria to end stage renal disease (ESRD) through a intermediate stage marked by microalbuminuria and overt proteinuria.^{1,2} Appearance of proteinuria heralds the onset of nephropathy and plays a critical role in the development of ESRD.³ So, quantifying urine protein accurately and precisely, is vital in the monitoring disease activity in patients with DM. The measurement of protein in 24-hr urine collection has been regarded as the gold standard.⁴ The use of 24-hr collection is necessitated by the variation in protein excretion throughout day which negates the use of concentration measurement in random urine collection.^{5,6} However, the 24-h urine collection is cumbersome, inconvenient especially in female patients, is often incomplete and difficult to administer in outpatients. A delay of 24-hr before definitive diagnosis further adds to the cost of the patients.⁷⁻¹¹ Some authors have even reported that 24-hr urine collection test is unreliable in up to one-third of cases due to incomplete urine collection.^{12,13} Unfortunately the other most widely used screening test for proteinuria-the dipstick test, has also been found to be fraught with error and correlates poorly with 24-h urine protein excretion.¹⁴⁻¹⁷

There remains therefore the need for a reliable quantitative measurement of urinary protein excretion that will be quick, easy to administer and correlate well with 24-hr urine protein excretion. The use of the urinary protein to creatinine ratio has been extensively demonstrated to possess the potential to fill this vacuum. But no consensus for specific Protein-creatinine (P:C) cutoff value has been obtained. These uncertainties have added to the resistance of substituting it for timed collections when evaluating diabetic nephropathy.

This study aims to evaluate the diagnostic value of P:C ratio in single voided urine samples for detection of proteinuria compared to those of a 24-hr samples in patients with diabetic nephropathy and also to determine the optimal cutoff for P:C ratio with best sensitivity and specificity for prediction of significant proteinuria.

MATERIALS AND METHODS

One hundred forty four patients aged 30-70, attending a Kantipur hospital who were diagnosed as diabetic were recruited in this study. The diabetic status was defined as per the American Diabetes Association (ADA).¹⁸ The study was undertaken for six months, from September 2008 to February 2009. The demographic data of patients including age, sex, body weight and height were obtained. 24-hr urine was collected from all participants. Participants were also instructed to collect un-timed spot

urine on the next day after the 24-hr urine collection. Informed consent was taken from all participants. Urine samples were collected at room temperatures without adding any preservatives. No specific recommendations were made about fluid intake, physical exercise or dietary protein intake. Women during menstruation period were not recruited for this study.

The adequacy of the 24-hr urine collection was assessed by comparing the total urinary creatinine in the sample with the predicted creatinine.

The predicted creatinine (mg/day) was calculated by- $[28 - (0.2 \times \text{age in years})] \times \text{weight in kg}$ for men and $[23.8 - (0.17 \times \text{age in years})] \times \text{weight in kg}$ for women. If the (predicted creatinine - measured creatinine)/predicted creatinine is >0.2 , the sample was considered as incomplete collection and was excluded from analysis.¹⁹

The 24-hr urine was stirred to ensure homogeneity and volume was measured in graduated cylinder and 1 ml aliquot sample was obtained. Urine creatinine was measured by modified Jaffe's method²⁰ [accurex, biomedical]. Urine protein was measured by pyrogallol red modified fujita method²¹ [chronolab sys S.L.]. All assays were performed, strictly adhering to standard operating procedures. The spot urine protein to creatinine ratio was obtained by dividing the urinary protein concentration by the urine creatinine concentration both expressed in mg/dl.

Statistical analysis: Statistical Package for Social Service (SPSS for window version; SPSS, 11.5, Inc., Chicago, IL) was used for statistical analysis. Spearman's rank correlation coefficient was used to show correlation between the spot urine p:c ratio and 24-hr urine total protein. Sensitivity, specificity and




Fig.1. Scatter plot between 24 hrs urinary protein excretion and Spot urine protein creatinine ratio ($r=0.892$)

predictive values of the random urine P:C ratios at various cutoffs for prediction of significant proteinuria were estimated using the results from the 24-hr urine protein as the gold standard. A Receiver operating characteristics (ROC) curve was plotted and area under the curve was calculated.

RESULTS

Among 144 subjects, 16 (11.5%) of urine samples were considered to be inadequately collected so excluded for analysis. In remaining 128 diabetic subjects (103 male and 25 female), 38 (29.0%) were found to have significant proteinuria by the gold standard, 24-hr urine protein estimation. A very good correlation was seen between the 24-hr urine protein and spot P:C ratio, with the Spearman's correlation (r) of 0.892 (Fig. 1).

The area under the ROC curve for random urine P:C ratio at various cutoffs is 0.88 (95.0% CI; 0.80-0.98 $p < 0.001$) (Fig. 2). Sensitivity and specificity of P:C ratio to detect proteinuria at various cutoffs is shown in Table-1. An excellent sensitivity of 95.6% and specificity of 74.5% were achieved to detect proteinuria at the P:C ratio cutoff greater than 0.15. With this cutoff, the positive predictive value was 61.1% and negative predictive value was 97.6%. At a cutoff 2.43, the specificity was 100.0% however the sensitivity drops to 52.2%, whereas at cutoff 0.07 the sensitivity became 100.0% limiting specificity only to 5.4%.

DISCUSSION

This study was conducted to evaluate the correlation between 24-hr urine protein and random urinary P:C ratio and to find the appropriate P:C cutoff for the prediction of significant proteinuria in diabetic subjects. Although some investigators advocate the use of albumin as an alternative to the total protein measurement²²⁻²⁴ and others have suggested that the profile of protein

Table-1: Sensitivity and Specificity of the P:C ratio at various cutoffs

Cutoff (mg/mg)	Sensitivity (%)	Specificity (%)	PV + *	PV - **
0.12	95.6	45.4	42.3	96.1
0.13	95.6	58.2	48.9	96.8
0.14	95.6	69.1	56.4	97.4
0.15	95.6	74.5	61.1	97.6
0.17	91.3	74.5	60.0	95.3
0.19	86.9	74.5	58.8	93.2
0.02	82.6	74.5	57.6	91.1
0.22	82.6	76.4	59.4	91.3

*Predictive value of positive test, **Predictive value of negative test

Table-2: Correlation of protein: creatinine ratio with 24-hr urine protein excretion shown in previous studies.

Author, year (Ref)	Correlation coefficient (r)
Ginsberg <i>et al</i> , 1983 ¹⁰	0.97
Houser <i>et al</i> , 1984 ²⁷	0.98
	0.97(ambulatory patient)
Schwab <i>et al</i> , 1987 ²⁸	0.98(hospitalized patient)
Lemann and Dumas, 1987 ²⁹	0.97
Boler <i>et al</i> , 1987 ⁹	0.85(Normal pregnancy)
	0.95(Twin pregnancy)
	0.96(hypertensive pregnancy)
Ralston <i>et al</i> , 1988 ³⁰	0.92
Jachevatsky <i>et al</i> , 1990 ³¹	0.92
Abitbol <i>et al</i> , 1990 ³²	0.95
Combs <i>et al</i> , 1991 ³³	0.98
Dyson <i>et al</i> , 1992 ³⁴	0.77
Mitchell <i>et al</i> , 1993 ³⁵	0.98
Quadri <i>et al</i> , 1994 ³⁶	0.92
Steinhauslin <i>et al</i> , 1995 ³⁷	0.93
Young <i>et al</i> , 1996 ³⁸	0.8
Robert <i>et al</i> , 1997 ³⁹	0.94
Sudan <i>et al</i> , 1997 ¹⁷	0.93
Ramos <i>et al</i> , 1998 ⁴⁰	0.94
Evans <i>et al</i> , 2000 ⁴¹	0.95
Rodriguez Thompson <i>et al</i> , 2001 ⁴²	0.8
Chitalia <i>et al</i> , 2001 ⁴³	0.97
Tornng <i>et al</i> , 2001 ⁴⁴	0.79
Yamasit <i>et al</i> , 2004 ⁴⁵	0.95
Leung YY <i>et al</i> , 2006 ⁴⁶	0.91
Alfredo <i>et al</i> , 2007 ⁴⁷	0.98
Wahbeh <i>et al</i> , 2009 ⁴⁸	0.83

excreted has differential diagnostic and prognostic Value, ²⁵ the National kidney foundation has recommended that an increased in protein excretion be used as a screening tools in patients at risk of developing renal disease. ²⁶ Since a rapid and accurate test avoid the inconvenience of patients as well as delay in diagnosis, a spot P: C ratio was taken as a rapid tool in our study to correlate it with 24-hr protein.

A good correlation was seen (r =0.892) in our study like in many other previous studies as shown in Table-2.

The excretion of creatinine and protein is reasonably constant throughout the day when the glomerular




Fig.2. ROC curve at various cutoffs for the spot random urinary protein-creatinine ratio (The area under the curve was 0.88 (95% confidence interval 0.80–0.98; P<0.001)

filtration rate is stable. ¹⁰ The urine P:C ratio corrects for variations in urinary protein concentration due to hydration and is not affected by a decrease in urine output in patients with renal insufficiency. So, the numerical outcome of the urine P:C ratio in mg/mg is roughly equal to the 24-hr protein excretion in g/day/1.73 m² body surface area.

Though our results agree with most of the authors, but still there are some well known conflicting results. Some authors have reported only a medium correlation as shown in Table-3. These discrepancies are probably due to an increase in tubular secretion of creatinine in various rates in patients with established renal disease ⁵³ and the interference of ketone bodies and some drugs in different creatinine assay method ⁵⁴ adopted by various authors. We have taken the P: C cutoff value of 0.15 to find out the correlation. Similar cutoff was taken in the study of Young *et al* ³⁸ and Mitchell *et al* ³⁵. The reference method cut off in (the study of Mitchell *et al* was taken as 150 mg/day whereas in the study of Young *et al* the reference method cutoff was taken as 300 mg/day.

We have taken the reference method cutoff as 150 mg/day. Because of the variability in laboratory methods for measuring proteinuria in different reported studies,

Table-3: Medium correlation shown between spot urine P:C ratio and 24-hr urinary protein excretion in previous studies

Author, year (Ref)	Correlation coefficient (r)
Lindow and Davey, 1992 ⁴⁹	0.53
Al <i>et al</i> , 2004 ⁵⁰	0.56
Durnwald and Mercer, 2003 ⁵¹	0.64
Aggarwal N <i>et al</i> , 2008 ⁵²	0.596

Table 4: Various P:C ratio cutoff and reference method cutoff taken in previous studies.

Author, year (Ref)	Patient group	Reference method cutoff (mg/day)	P:C ratio cutoff (mg/mg)
Ginsberg <i>et al</i> , 1983 ¹⁰	Renal clinic	200	0.2
Ralston <i>et al</i> , 1988 ³⁰	Rheumatology clinic	300	0.35
Dyson <i>et al</i> , 1992 ³⁴	Renal transplant clinic	500	0.35
Mitchell <i>et al</i> , 1993 ³⁵	Elderly patient clinic	150	0.15
Quadri <i>et al</i> , 1994 ³⁶	Pregnant patient	300	0.3
Young <i>et al</i> , 1996 ³⁸	Pregnant patient	300	0.15
Robert <i>et al</i> , 1997 ³⁹	Pregnant patient	300	0.17
Sudan <i>et al</i> , 1997 ¹⁷	Pregnant patient	300	0.26
Ramos <i>et al</i> , 1998 ⁴⁰	Pregnant patient	300	0.5
Evans <i>et al</i> , 2000 ⁴¹	Pregnant patient	300	0.3
Rodriguez <i>et al</i> , 2001 ⁴²	Pregnant patient	300	0.19
Chitalia <i>et al</i> , 2001 ⁴³	Renal clinic	250	0.26
Tornig <i>et al</i> , 2001 ⁴⁴	Renal transplant clinic	500	0.35
Durnwald and Mercer, 2003 ⁵¹	Pregnant patient	300	0.3
Yamasit <i>et al</i> , 2004 ⁴⁵	Pregnant patient	300	0.19
Al <i>et al</i> , 2004 ⁵⁰	Pregnant patient	300	0.19

several cutoff points and different units for the urinary P:C ratio have been reported thereby precluding valid comparisons among such studies. The discrepancies shown above in the value of correlation coefficient also might be due to this heterogeneity of cutoff value adopted in different studies. reported studies, several cutoff points and different units for the urinary P:C ratio have been reported thereby precluding valid comparisons among such studies. The discrepancies shown above in the value of correlation coefficient also might be due to this heterogeneity of cutoff value adopted in different studies.

Not only the P:C ratio cutoff variation is seen in various studies but also the variations in reference method cutoff have been noted (Table-4).

In our study the ROC curve analysis showed an area under the curve of 0.88 indicating that the urinary P:C ratio is sufficiently accurate to predict significant proteinuria. At the optimal cutoff point of 0.15, the diagnostic performance was high with sensitivity and specificity of 96.65 and 74.4% respectively. The higher value found for sensitivity compared with specificity would suggest that the ratio test might be more valuable as a rule out test. Similarly the higher negative predictive value of 97.6% in comparison to positive predictive value of only 61.1% supports this tentative conclusion.

In conclusion, random P:C ratio is a good predictor of significant proteinuria in diabetic nephropathy. Since various studies have given their own reference method and P:C ratio cutoff , it is necessary to find out different cutoff (having optimal sensitivity and specificity) for different group of patients in our own laboratory settings to replace the gold standard method.

REFERENCES

- Selby JV, Fitzmonns SC, Newman JM, Kats PP, Sepe S, Showstack J. The natural history of diabetic nephropathy. *J Amer Med Assoc* 1990; 263: 1945-60.
- Andrzej SK, James HW. Natural history of diabetic nephropathy. *Diabetes Rev* 1995; 3: 446-57.
- Campbell RC, Ruggenenti P, Remuzzi G. Proteinuria in diabetic nephropathy:Treatment and Evolution. *Curr Diab Rep* 2003; 3: 497-504.
- Price CP, Newall RG, Boyd JC. Use of protein: Creatinine ratio measurements on random urine samples for prediction of significant proteinuria: A systemic review. *Clin Chem* 2005; 51(9): 1577-86.
- Koopman MG, Krediet RT, Koomen GM, Strackee J, Arisz L. Circadian rhythm of proteinuria: Consequences of the use of protein:creatinine ratios. *Nephrol Dial Transplant* 1989; 4: 9-14.
- Kassirer JP, Harrington JT. Laboratory evaluation of renal function. In: Schrier RW, Gottschalk CW, editors. *Diseases of the kidney* (4th ed.). Vol 1. Little Brown Boston 1988. 393-426.
- Shaw AB, Risdon P, lewis-Jackson JD. Protein creatinine index and albustix in assessment of proteinuria. *Brit Med J* 1983; 287: 929-32.

8. Ruggenenti P, Gaspari F, Perna A, Remuzzi G. Cross sectional longitudinal study of spot morning urine protein:creatinine ratio, 24-hr urine protein excretion rate, glomerular filtration rate, and end stage renal failure in chronic renal disease in patients without diabetes. *Brit Med J* 1998; 316: 504-9.
9. Boler L, Zbella EA, Gleicher N. Quantitation of proteinuria in pregnancy by the use of single voided urine samples. *Obstet Gynecol* 1987; 70: 99-100.
10. Ginsberg JM, Chang BS, Matarese RA, Garella S. Use of single voided urine samples to estimate quantitative proteinuria. *N Engl J Med* 1983; 309: 1543-6.
11. Austin HA. Clinical evaluation and monitoring of lupus kidney disease. *Lupus* 1998; 7: 618-21.
12. Christopher SL, Petri M, Astor BC, Fine D. Urine protein-to-creatinine ratio is a reliable measure of proteinuria in lupus nephritis. *J Rheumatol* 2004; 31: 1557-9.
13. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of diet in renal disease study group. *Ann Intern Med* 1999; 130: 461-70.
14. Uttendorfsky OT, Veersema D, Greene T, Mooij PM, Stolte LM. Protein/creatinine ratio in the assessment of proteinuria during pregnancy. *Eur J Obstet Gynecol Repro Biol* 1988; 27: 221-26.
15. Sibai BM. Pitfalls in diagnosis and management of pre-eclampsia. *Amer J Obstet Gynecol* 1998; 159: 1-5.
16. Irgensmoller L, Hemmingsen L, Holm J. Diagnostic value of microalbuminuria in pre-eclampsia. *Clin Chem Acta* 1986; 158: 295-8.
17. Saudan PJ, Brown MA, Farrel T, Shaw L. Improved methods in assessing proteinuria in hypertensive pregnancy. *Br J Obstet Gynecol* 1997; 104: 1159-64.
18. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: follow-up report on the diagnosis of diabetes mellitus. *Diabetes Care* 2003; 26: 3160-67.
19. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron* 1976; 16: 31-41.
20. Chasson AL, Crady HJ, Stanley MA. Determination of creatinine by means of automated chemical analysis. *Amer J Clin Pathol* 1961; 35: 83-8.
21. Chambers RE, Bullock DG, Whicher JT. External quality assessment of total urinary protein estimation in the United Kingdom. *Ann Clin Biochem* 1991; 28: 467-73.
22. Ballantyne FC, Gibbon J, O'reilly D. Urine albumin should replace total protein for the assessment of glomerular proteinuria. *Ann Clin Biochem* 1993; 30: 101-3.
23. Beetham R, Cattell WR. Proteinuria: Pathophysiology, significance and recommendations in clinical practice. *Ann Clin Biochem* 1993; 30: 425-34.
24. Newman DJ, Thakkar H, Medcalf EA, Gray MR, Price CP. Use of urine albumin measurement as a replacement for total proteinuria. *Clin Nephrol* 1995; 43: 104-9.
25. Hofmann W, Guder WG. A diagnostic programme for quantitative analysis of proteinuria. *J Clin Chem Biochem* 1989; 27: 589-600.
26. National kidney foundation K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Amer J Kidney Dis* 2002; 39(2 suppl 1): S1-S266.
27. Houser M. Assessment of proteinuria using random urine samples. *J Pediatr* 1984; 104: 845-8.
28. Schwab SJ, Christensen L, Dougherty K, Klahr S. Quantification of proteinuria by the use of protein-to-creatinine ratios in single urine samples. *Arch Intern Med* 1987; 147: 943-4.
29. Lemann J, Doumas B. Proteinuria in health and disease assessed by measuring the urinary protein/ creatinine ratio. *Clin Chem* 1987; 33: 297-9.
30. Ralston SH, Caine N, Richards I, O'Reilly D, Sturrock RD, Capell HA. Screening for proteinuria in a rheumatology clinic: comparison of dipstick testing, 24-hr urine quantitative protein and protein/creatinine ratio in random urine samples. *Ann Rheum Dis* 1998; 47: 759-63.
31. Jachevastzky OE, Rosenberg RP, Shalit A, Zonder HB, Grunstein S. Protein/creatinine ratio in random urine specimens for quantification of proteinuria in preeclampsia. *Obstet Gynecol* 1990; 75: 604-6.
32. Abitbol C, Zilleruelo G, Freundlich M, Strauss J. Quantitation of proteinuria with urinary protein /creatinine ratios and random testing with dipsticks in nephritic children. *J Pediatr* 1990; 116: 243-7.
33. Combs CA, Wheeler BC, Kitzmiller JL. Urinary protein/creatinine ratio before and during pregnancy in women with diabetes mellitus. *Amer J Obstet Gynecol* 1991; 165: 920-3.
34. Dyson EH, Will EJ, Davison AM, O'Malley AH, Shepherd HT, Jones RG. Use of the urinary protein creatinine index to assess proteinuria in renal transplant patients. *Nephrol Dial Transplant* 1992; 7: 450-2.
35. Mitchell SM, Sheldon TA, Shaw AB. Quantification of proteinuria: A reevaluation of the protein/creatinine ratio for elderly subjects. *Age Aging* 1993; 22: 443-9.
36. Quadri KM, Bernardini J, Greenberg A, laifer S, Syed A, Holley JL. Assessment of renal function during pregnancy using random urine protein to creatinine ratio and cockcroft-Gault formula. *Amer J kidney Dis* 1994; 24: 416-20.
37. Steinhauslin F, Wauters JP. Quantitation of proteinuria in kidney transplant patients: accuracy of the urine protein/creatinine ratio. *Clin Nephrol* 1995; 43: 110-15.
38. Young RA, Buchanan RJ, Kinch RA. Use of the protein/creatinine ratio of a single voided urine specimen in the evaluation of suspected pregnancy-induced hypertension. *J Fam Pract* 1996; 42: 385-9.
39. Robert M, Sepandj F, Liston RM, Dooley KC. Random protein-creatinine ratio for the quantitation of proteinuria in pregnancy. *Obstet Gynecol* 1997; 90: 893-5.
40. Ramos JGL, Martins-costa SH, Mathias MM, Guerin YLS, Barros EG. Urinary protein/creatinine ratio in hypertensive pregnant women. *Hypertens Pregnancy* 1999; 18: 209-18.
41. Evans W, Lensmeyer JP, Kirby RS, Malnory ME, Broekhuizen FF. Two hour urine collection for evaluating renal function correlates with 24-hr urine collection in pregnant patients. *J Matern Fetal Med* 2009; 9: 233-7.
42. Rodriguez-Thompson D, Lieberman ES. Use of a random urinary protein to creatinine ratio for the diagnosis of significant proteinuria during pregnancy. *Amer J Obstet Gynecol* 2001; 185: 808-11.
43. Chitalia VC, Kothari J, Wells EJ, et al. Cost benefit analysis and prediction of 24-hr proteinuria from the spot urine protein creatinine ratio. *Clin Nephrol* 2001; 55: 436-47.
44. Torng S, Rigatto C, Rush DN, Nickerson P, Jeffery JR. The urine protein to creatinine ratio as a predictor of 24-hr urine protein excretion in renal transplant patients. *Transplant* 2001; 72: 1453-6.
45. Yamasit W, Chaithongwongwatthana S, Charoenvidhya D,

- Uerpairajkit B, Tolosa J. Random urinary protein to creatinine ratio for prediction of significant proteinuria in women with preeclampsia. *J Matern Fetal Neonatal Med* 2004; 16: 275-9.
46. Leung YY, Szeto CC, Tam LS, et al. Urine protein to creatinine ratio in an untimed urine collection is a reliable measure of proteinuria in lupus nephritis. *Rheumatology* 2007; 46: 649-52.
47. Alfredo LM, Jeneth MA, Fernando RA, et al . Protein creatinine ratio in random urine samples is a reliable marker of increased 24-hr protein excretion in hospitalized women with hypertensive disorders of pregnancy. *Clin Chem* 2007; 53(9): 1623-8.
48. Wahbeh MA, Ewais HM, Elsharif ME. Comparison of 24-hr urinary protein and protein to creatinine ratio in the assessment of proteinuria. *Saudi J kidney Dis Transpl* 2009; 20: 443-7.
49. Lindow SW, Davey DA. The variability of urinary protein and creatinine excretion in patients with gestational proteinuric hypertension. *Brit J Obstet Gynaecol* 1992; 99: 869-72.
50. Al RA, Baykal C, Karacay O, Geyik PO, Altun S, Dolen L. Random urine protein creatinine ratio to predict proteinuria in new-onset mild hypertension in late pregnancy. *Obstet Gynecol* 2004; 104: 367-71.
51. Durnwald C, Mercer B. A prospective comparison of total protein/creatinine ratio versus 24-hr urine protein in women with suspected pre-eclampsia. *Amer J Obstet Gynecol* 2003; 189: 848-52.
52. Aggarwal N, Suri V, Soni S, Kohli H. A prospective comparison of random urine protein-creatinine ratio vs 24-hr urine protein in women with preeclampsia. *Medscape J Med* 2008; 10: 98-102.
53. Carrie BJ, Golbetz HV, Michaelis AS, Myers BD. Creatinine: An inadequate filtration marker in glomerular disease. *Amer J Med* 1980; 60: 177-82.
54. Payne RB. Creatinine clearance: a redundant clinical investigation. *Ann Clin Biochem* 1986; 23: 243-50.