

Association of thyroid dysfunction among infertile women visiting infertility center of Om Hospital, Kathmandu, Nepal

B Rijal,¹ R Shrestha² and B Jha¹

¹Om Hospital and Research Centre, Kathmandu, Nepal, ²Department of Biochemistry, Nepal Medical College and Teaching Hospital, Kathmandu, Nepal

Corresponding author: Dr. Bhola Rijal, Om Hospital and Research Centre, Kathmandu, Nepal

ABSTRACT

Infertility is one of the medial, social and psychological burdens in this part of world. Thyroid dysfunction can lead to menstrual disturbance, anovulatory cycles, and decreased fecundity. Proper management of thyroid dysfunction can result restoration of normal fertility. Therefore it is very important to screen thyroid abnormalities among women with infertility. This study aimed to determine association of thyroid dysfunction among infertile women. This study comprises total of 735 primary infertile women with age ranging from 20 to 35 years. Blood samples were collected and subjected for estimation of thyroid hormones. Out of 735 cases 547 (74.4%) were euthyroid. 56 (7.6%) have primary hypothyroidism, 31 (4.2%) have primary hyperthyroidism, 87 (11.8%) have subclinical hypothyroidism, and 8 (1.1%) have subclinical hyperthyroidism.

Keywords: Infertility, Thyroid hormone, Thyroid dysfunction

INTRODUCTION

Infertility is defined as the inability to conceive after one year of regular intercourse without contraception. It is one of the medial, social and psychological burden in this part of world. About 18–20% of couples in reproductive age are infertile.¹ Endocrines play central role in maintaining fertility. Thyroid hormones have profound effects on reproduction and pregnancy. There is a known association of hyperthyroidism and hypothyroidism with menstrual disturbances and decreased fecundity. Therefore assessment of thyroid dysfunction has been considered as an important component of infertility work up in women.² There are several experimental evidences, both on animals and humans, indicating association of hyperthyroidism or hypothyroidism with menstrual disturbance, anovulatory cycles, decreased fecundity and increased morbidity during pregnancy.³⁻⁵ Most importantly proper management of such thyroid dysfunction results in improvement in health status, normalization of menstrual abnormalities and restoration of normal fertility.⁶ Therefore it is very important to screen thyroid abnormalities among women with infertility, particularly in countries like us, considered as areas with endemic goiter. Infertility associated with thyroid dysfunction in these areas is not uncommon.⁷

The aim of this study was to find association of thyroid dysfunction among infertile women visiting our infertility and IVF center.

MATERIALS AND METHODS

This study comprises total of 735 primary infertile women with age ranging from 20 to 35 years, visiting infertility and IVF center of Om Hospital and Research

Centre, Kathmandu, Nepal. The cases were selected over period of two years (March 2009 to 2011). The inclusion criteria for the selection of cases were diagnosis of primary infertility, age between 20-35 years with duration of marriage more than one year. Infertility due to male factor and the female factors with any congenital anomaly of urogenital tract, or any obvious organic lesion were excluded. Any history of thyroid disease, thyroid surgery and those treated with thyroid medication were also excluded.

Five milliliters of venous blood was collected in fasting state. Serum was prepared within one hour of blood collection and stored at -20°C till get analyzed for thyroid function test. All samples were analyzed within 15 days from blood collection date. Serum free 3,5,3'-triiodothyronine (fT3), free 3,5,3',5'-tetraiodothyronine (fT4) and thyroid-stimulating hormone (TSH) were analyzed using VitrosECi™ analyzer (Ortho Clinical Diagnostics; OCD, Rochester, NY). Measurement of fT3 and fT4 was based on a direct, labeled antibody, competitive immunoassay, whereas immunometric immunoassay technique was used in measurement of TSH. The bound HRP conjugate was measured by a luminescent reaction.⁸ Assay validity and reliability was determined by use of control sera provided by OCD. Normal range used for the third generation TSH assay was 0.46 to 4.5 iIU/mL, and cut off range for fT3 and fT4 used were 4.26 to 8.10 pmol/L and 10.0 to 28.2 pmol/L respectively.

As per thyroid profile cases were categorized into 7 groups- Euthyroidism, primary hypothyroidism, primary hyperthyroidism, secondary hypothyroidism, secondary hyperthyroidism, subclinical hypothyroidism, and

Table-1: Categorization of cases on the basis of thyroid profile

	ft3 (pmol/L)	ft4 (pmol/L)	TSH (?IU/mL)
Euthyroid	4.26 - 8.10	10.0 - 28.2	0.46 - 4.5
Primary Hypothyroid	< 4.26	< 10.0	> 4.5
Primary Hyperthyroid	> 8.1	> 28.2	< 0.46
Secondary Hypothyroid	< 4.26	< 10.0	< 0.46
Secondary Hyperthyroid	> 8.1	> 28.2	> 4.5
Subclinical Hypothyroid	4.26 - 8.10	10.0 - 28.2	> 4.5
Subclinical Hyperthyroid	4.26 - 8.10	10.0 - 28.2	< 0.46

subclinical hyperthyroidism (Table-1).⁹⁻¹⁴ The statistical analysis was carried out using Statistical Package for Social Sciences SPSS 14 software for windows.

RESULTS

The thyroid function status of our study population was elucidated in Table-2. Mean age of the study subjects were 26.3± 3.1 years. There was no significant difference in age among different groups according to thyroid status (p>0.05). Out of 735 cases 547 (74.4%) were euthyroid. Large proportion of the cases that is 25.6% of the cases have different types of thyroid dysfunction (p<0.05). Among cases with thyroid dysfunction, 56 (7.6%) have primary hypothyroidism, 31 (4.2%) have primary hyperthyroidism, 87 (11.8%) have subclinical hypothyroidism, and 8 (1.1%) have subclinical hyperthyroidism. Only 5 (0.7%) and 1 (0.1%) have secondary hypothyroidism and secondary hyperthyroidism respectively.

DISCUSSION

This study revealed that 25.6% of our study populations have thyroid dysfunction and possibly resulted infertility. Majority of thyroid dysfunction was hypothyroidism comprising almost 20.1%, out of which subclinical hypothyroidism being most common (11.8%).

Subclinical hypothyroidism appears to one of the important thyroid dysfunction resulting infertility.¹⁵ Subclinical hypothyroidism may be associated with ovulatory dysfunction and adverse pregnancy outcome. Our results of high prevalence of hypothyroidism in infertile women is also compatible with study conducted in our neighborhood countries.^{16,17} Higher prevalence of hypothyroidism may be precipitated due to higher prevalence of iodine deficiency in this country, as it lies in iodine deficient range.¹⁸ Overall prevalence of iodine deficiency in Nepal is 13.6%.¹⁹ Prevalence of hypothyroidism is even high in pregnant women in compare to non-pregnant women as demonstrated by one of study conducted in Nepal.²⁰ Hypothyroidism due to chronic autoimmune thyroiditis may also contribute significantly. Relatively higher prevalence rate of hypothyroidism in our study can be also due to special referral pattern of the patients who were referred to the hospital based on suspicion of thyroid abnormalities.

In women of fertile age, hypothyroidism results oligomenorrhea and amenorrhea, polymenorrhea, and menorrhagia. High frequency of infertility in hypothyroidism maybe due to altered peripheral estrogen metabolism, hyperprolactinemia, and disturbances in gonadotropin releasing hormone (GnRH) secretion that result in an abnormal pulsatile release of LH.²¹ Moreover, thyroid hormones appear necessary to achieve maximum fertilization rates and have special role in oocyte physiology. Furthermore, serum TSH levels can be a significant predictor of fertilization failure in women undergoing IVF.²² Most importantly, treatment of hypothyroidism with thyroid drugs has been shown to normalize prolactin levels as well as normal LH response to LHRH, reduce menstrual disturbances, and finally increase the chances of spontaneous fertility.^{23,24} Although menstrual irregularities are common, ovulation and conception can still occur in hypothyroidism, where

thyroxine treatment restores a normal menstrual pattern and reverses hormonal changes. Therefore, it is very important to identify the hypothyroidism in infertility.

Overall prevalence of hyperthyroidism in our study population is 5.4%, in which primary hyperthyroidism is most common. Another study conducted in India in similar setting also have similar prevalence of hyperthyroidism in

Table-2: Distribution of cases according to thyroid status and mean of thyroid hormone

	N (735)	%	hormone		
			ft3 (pmol/L) Mean ± SD	ft4 (pmol/L) Mean ± SD	TSH (?IU/mL) Mean ± SD
Euthyroid	547	74.4	5.3 ± 1.5	15.1 ± 3.1	2.1 ± 1.6
Primary Hypothyroid	56	7.6	2.1 ± 1.1	4.2 ± 2.5	10.3 ± 5.2
Primary Hyperthyroid	31	4.2	14 ± 1.8	36 ± 3.3	0.2 ± 0.1
Secondary Hypothyroid	5	0.7	2.2 ± 1.0	5.1 ± 2.0	0.3 ± 0.1
Secondary Hyperthyroid	1	0.1	15 ± 1.2	33 ± 1.8	9.1 ± 1.5
Subclinical Hypothyroid	87	11.8	5.5 ± 1.4	14.6 ± 2.8	9.5 ± 1.2
Subclinical Hyperthyroid	8	1.1	5.2 ± 1.3	16.0 ± 2.1	0.3 ± 0.1

infertile women.²⁵ Thyrotoxicosis results change in serum sex hormone binding globulin, thus result changes in sex hormone concentration. Changes in sex hormone concentration including LH and FSH, result menstruation disturbance including oligomenorrhea, hypomenorrhea, and anovulation.²⁶⁻²⁹ menstrual disturbances, frequent in thyrotoxicosis are restored following treatment.

One of the major drawback of this study was other possible hormonal causes of infertility was not investigated. Specific pattern of thyroid dysfunction in our study patients were not analyzed.

There is high prevalence of thyroid dysfunction among the infertile women. Proper management of the thyroid dysfunction can result regain of fertility. Therefore, routine screening is required to all cases of infertility for possible thyroid disorders.

REFERENCE

- Putowski L, Darmochwal-Kolarz D, Rolinski J, Oleszczuk J, Jakowicki J. The immunological profile of infertile women after repeated IVF failure (Preliminary study). *Eur J Obstet Gynecol Reprod Biol* 2004; 112: 192-6.
- Cramer DW, Sluss PM, Powers RD *et al*. Serum prolactin and TSH in an in vitro fertilization population: is there a link between fertilization and thyroid function? *J Assist Reprod Genet* 2003; 20: 210-5.
- Poppe K, Velkeniers B. Thyroid disorders in infertile women. *Ann Endocrinol (Paris)* 2003; 64: 45-50.
- Doufas AG, Mastorakos G. The hypothalamic-pituitary-thyroid axis and the female reproductive system. *Ann N Y Acad Sci* 2000; 900: 65-76.
- Poppe K, Velkeniers B, Glinoer D. Thyroid disease and female reproduction. *Clin Endocrinol (Oxf)* 2007; 66: 309-21.
- Miciński P, Wielgus E, Wojcieszyn M, Pawlicki K. Abnormal ovarian reserve test reflects thyroid dysfunction. *Poland J Gyn Invest* 2006; 9: 30-4.
- Zimmermann MB, Jooste PL, Pandav CS. Iodine-deficiency disorders. *Lancet* 2008; 372: 1251-62.
- Summers M. Luminogenic Reagent Using 3-Chloro 4-Hydroxy Acetanilide to Enhance Peroxidase/Luminol Chemiluminescence. *Clin Chem* 1995; 41: S73.
- Evered D. Diseases of the Thyroid Gland. *Clin Endocrinol Metabol* 1974; 3: 425-50.
- Evered DC, Ormston BJ, Smith PA *et al*. Grades of hypothyroidism. *Brit Med J* 1973; 1: 657-62.
- Caldwell G, Gow SM, Sweeting VM *et al*. A new strategy for thyroid function testing. *Lancet* 1985; 1: 1117-9.
- Kerr DJ, Alexander WD. Is the TRH test usually unnecessary? *Lancet* 1984; 2: 1161-2.
- Spencer CA. Thyroid profiling for the 1990's: free T4 estimate or sensitive TSH measurement. *J Clin Immunol* 1989; 12: 82-9.
- Nicoloff JT, Spencer CA. The use and misuse of the sensitive thyrotropin assays. *J Clin Endocrinol Metabol* 1990; 71: 553-8.
- Bohnet HG, Fiedler K, Leidenberger FA. Subclinical hypothyroidism and infertility. *Lancet* 1981; 2: 1278.
- Goswami B, Patel S, Chatterjee M, Koner BC, Saxena A. Correlation of Prolactin and Thyroid Hormone Concentration with Menstrual Patterns in Infertile Women. *J Reprod Infertile* 2009; 10: 207-12.
- Elahi S, Tasneem A, Nazir I, Nagra SA, Hyder SW. Thyroid dysfunction in infertile women. *J Coll Physicians Surg Pakistan* 2007; 17: 191-4.
- Pandav CS. IDD in Southeast Asia. In: Hetzel BS, Pandav CS, eds. SOS for a billion, the conquest of iodine deficiency disorders. Delhi: Oxford University Press, 1994: 213-31.
- Baral N, Ramprasad C, Lamsal M, Koner BC, Koirala S. Assay of iodine deficiency status in three ecological regions of Nepal by a microdigestion method. *Southeast Asian J Trop Med Public Health* 1999; 30: 527-31.
- Khanal M, Malla S, Sharma V, Gyawali P, Shrestha M, Shrestha R. Assessment of thyroid function in pregnant subjects. *J Nepal Assoc Med Lab Sci* 2009; 10: 39-41.
- Krassas GE. Thyroid disease, menstrual function and fertility. *Thyroid International* 2000; 1: 1-15180
- Cramer DW, Sluss PM, Powers RD *et al*. Serum prolactin and TSH in an in vitro fertilization population: is there a link between fertilization and thyroid function? *J Assist Reprod Genet* 2003; 20: 210-5.
- Redmond GP. Thyroid dysfunction and women's reproductive health. *Thyroid* 2004; 14(Suppl 1): S5-S15.
- Krassas GE. Thyroid disease and female reproduction (modern trends). *Fertil Steril* 2000; 74: 1063-70.
- Joshi JV, Bhandarkar SD, Chadha M, Balaiah D, Shah R. Menstrual irregularities and lactation failure may precede thyroid dysfunction or goitre. *J Postgrad Med* 1993; 39: 137-41.
- Benson RC, Dailey ME. The menstrual pattern in hyperthyroidism and subsequent posttherapy hypothyroidism. *Surg Gynecol Obstet* 1955; 100: 19-26.
- RE, Sturgis SH, Lerman J, Stanbury JB. The menstrual pattern in thyroid disease. *J Clin Endocrinol Metabol* 1952; 12: 846-55.
- Joshi JV, Bhandarkar SD, Chadha M, Balaiah D, Shah R. Menstrual irregularities and lactation failure may precede thyroid dysfunction or goitre. *J Postgrad Med* 1993; 39: 137-41.
- Tanaka T, Tamai H, Kuma K, Matsuzuka F, Hidaka H. Gonadotropin response to luteinizing hormone releasing hormone in hyperthyroid patients with menstrual disturbances. *Metabol* 1981; 30: 323-6.